

**Supplementary Data. Shedding dynamics of Morogoro virus, an Old World
Arenavirus closely related to Lassa virus, in its natural reservoir host *Mastomys
natalensis***

**Benny Borremans,¹ Raphaël Vossen,¹ Beate Becker-Ziaja,² Sophie Gryseels,¹ Nelika Hughes,¹
Natalie Van Houtte,¹ Stephan Günther,² Herwig Leirs¹**

¹Evolutionary Ecology Group, University of Antwerp, B-2020 Antwerp, Belgium; and ²Bernhard-
Nocht-Institute for Tropical Medicine, D-20359 Hamburg, Germany

Inhibition test




Figure 1. Inhibition test results for blood from capillaries (cap) and filter paper (fp), saliva, urine and feces. The line shows the estimated standard curve as a reference.




Figure 2. Log-transformed antibody titers (blood) and Morogoro RNA (blood, excretions) for individual 1 of 15.




Figure 3. Log-transformed antibody titers (blood) and Morogoro RNA (blood, excretions) for individual 2 of 15.




Figure 4. Log-transformed antibody titers (blood) and Morogoro RNA (blood, excretions) for individual 3 of 15.




Figure 5. Log-transformed antibody titers (blood) and Morogoro RNA (blood, excretions) for individual 4 of 15.




Figure 6. Log-transformed antibody titers (blood) and Morogoro RNA (blood, excretions) for individual 5 of 15.




Figure 7. Log-transformed antibody titers (blood) and Morogoro RNA (blood, excretions) for individual 6 of 15.




Figure 8. Log-transformed antibody titers (blood) and Morogoro RNA (blood, excretions) for individual 7 of 15.




Figure 9. Log-transformed antibody titers (blood) and Morogoro RNA (blood, excretions) for individual 8 of 15.




Figure 10. Log-transformed antibody titers (blood) and Morogoro RNA (blood, excretions) for individual 9 of 15. This individual died on day 12 post-infection.




Figure 11. Log-transformed antibody titers (blood) and Morogoro RNA (blood, excretions) for individual 10 of 15.




Figure 12. Log-transformed antibody titers (blood) and Morogoro RNA (blood, excretions) for individual 11 of 15.




Figure 13. Log-transformed antibody titers (blood) and Morogoro RNA (blood, excretions) for individual 12 of 15.




Figure 14. Log-transformed antibody titers (blood) and Morogoro RNA (blood, excretions) for individual 13 of 15. This individual died on day 140 post-infection.




Figure 15. Log-transformed antibody titers (blood) and Morogoro RNA (blood, excretions) for individual 14 of 15.




Figure 16. Log-transformed antibody titers (blood) and Morogoro RNA (blood, excretions) for individual 15 of 15. This individual died on day 130 post-infection.